



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Hence

$$\binom{n}{a} - \binom{i}{1} \binom{n-1}{a} + \binom{i}{2} \binom{n-2}{a} - \cdots + (-1)^i \binom{n-i}{a} = \binom{n-i}{a-i}. \quad (1)$$

In the case when  $i = a$ , we have  $a - i = 0$ , and  $\binom{n-i}{a-i} = 1$ ; therefore, equation (1) becomes

$$\binom{n}{a} - \binom{a}{1} \binom{n-1}{a} + \binom{a}{2} \binom{n-2}{a} - \cdots + (-1)^a \binom{n-a}{a} = 1. \quad (2)$$

When  $i > a$ , then  $a - i$  is negative, so that  $\binom{n-i}{a-i} = 0$ . Hence for  $i > a$ ,

$$\binom{n}{a} - \binom{i}{1} \binom{n-1}{a} + \binom{i}{2} \binom{n-2}{a} - \cdots + (-1)^i \binom{n-i}{a} = 0. \quad (3)$$

An excellent solution by a somewhat different method from the ones exhibited in these two solutions was received from A. M. HARDING.

**425. Proposed by CLIFFORD N. MILLS, Brookings, S. D.**

Solve for  $x$  and  $y$  the equations  $2^{x+y} = 6$ ,  $2^{x+1} = 3^y$ .

SOLUTION BY ELIJAH SWIFT, University of Vermont.

We shall agree to mean by  $a^x$ , where  $a$  is real,  $e^{x \log a}$ , in which the real logarithm of  $a$  is to be taken, so that  $a^x$  is single valued. We have, then, from the given equations,  $(x+y) \log 2 = \log 6$ ,  $(x+1) \log 2 = y \log 3$ , where all the logarithms are real. Solving these linear equations for  $x$  and  $y$ , we obtain the required solutions

$$x = \frac{(\log 6)^2 - \log 2 \cdot \log 12}{\log 2 \log 6} = 1.198 \quad \text{and} \quad y = \frac{\log 12}{\log 2} = 1.387.$$

On account of the homogeneity of these fractions in the logarithmic function, we may substitute logarithms to the base 10 in our computation of the numerical values of  $x$  and  $y$ .

Also solved by EMMA GIBSON, NATHAN ALTSHILLER, A. L. MCCARTY, A. M. HARDING, HORACE OLSON, RICHARD MORRIS, G. W. HARTWELL, C. E. GITHENS, FRANK IRVIN, ELIZABETH B. DAVIS, F. L. CARMICHAEL, EDWARD S. INGHAM, S. A. JOFFE, W. C. EELLS, ELMER SCHUYLER, V. M. SPUNAR, R. M. MATHEWS, and CYRIL A. NELSON.

GEOMETRY.

**454. Proposed by LOUIS ROUILLION, Mechanics Institute, New York City.**

Show how to construct an equilateral triangle with its vertices lying on three [parallel] lines not equally spaced.

I. SOLUTION BY C. N. SCHMALL, New York City.

The proposer evidently meant that the three lines are parallel.

(i) Let  $x$ ,  $y$ , and  $z$  be the three parallel lines in order. At any point,  $M$ , on the line  $y$ , construct the angles  $AMC$  and  $BMC$  each equal to  $60^\circ$  (Fig. 1). Draw the line  $AB$ , and, about the triangle  $AMB$ , circumscribe the circle  $ABC$ .